

University of Wisconsin-Extension

GEOLOGICAL AND NATURAL HISTORY SURVEY
3817 Mineral Point Road
Madison, Wisconsin 53705

James Robertson, State Geologist and Director

DANE COUNTY ATRAZINE PROJECT DATABASE AND
DOCUMENTATION: PROGRAMS, DATA SOURCES AND
METHODS

by

M F. Bohn

Open-File Report 1994-01
3 p. + 10 3 1/2 inch high-density diskettes

This report represents work performed by the Geological and Natural History Survey, and is released to the open files in the interest of making the information more readily available. This report has not been edited or reviewed for conformity with Geological and Natural History Survey standards and nomenclature.

1994

Wisconsin Geological and Natural History Survey Open-file Report 1994-01

3 p + 10 3-1/2 inch high-density diskettes

Dane County Atrazine Project Database and Documentation: Programs, Data Sources and Methods

by

M.F. Bohn

This is the supporting programs, data and methods for WOFR 1994-02, Hydrogeologic and land-use controls on atrazine detections in Dane County Wisconsin by M.A. Bohn, F.W. Madison, M. Muldoon, and K. Connors. A copy of Summary for WOFR1994-02 is attached.

The information consists of 909 files in Pkzip compressed format totalling 12,402,194 bytes. The expanded files consist of various geology, hydrogeology, land use, topographic and geographic information in Arc format.

No documentation was provided.

Title: HYDROGEOLOGIC AND LAND-USE CONTROLS ON ATRAZINE DETECTIONS IN DANE COUNTY, WISCONSIN

Investigators: Michael A. Bohn, GIS Specialist; Frederick W. Madison, Soil Scientist; Maureen Muldoon, Hydrogeologist; Wisconsin Geological and Natural History Survey, University of Wisconsin-Extension. Kevin Connors, County Conservationist, Dane County Land Conservation Department.

Period of Contract: August 5, 1991 - June 30, 1993

Objectives: The primary objective of this project is to determine the soils, geologic, and hydrogeologic factors that affect atrazine contamination of domestic wells.

**Background/
Need:**

Atrazine, a popular and inexpensive herbicide, has been used extensively during the past 30 years in the corn-producing regions of the Upper Midwest. Concern over the possible movement of atrazine to groundwater led to state-wide sampling programs, site-specific investigations of atrazine movement, and approval of the 1991 Atrazine Rule which established atrazine management areas where the use of atrazine is restricted and prohibition areas.

The Rural Well Survey (LeMasters, 1990) indicated that 50% of the rural wells in Dane County, an area of dairy-livestock agriculture with high corn production, contained detectable levels of triazine-based compounds.

There was interest in trying to relate atrazine detections to land-use and hydrogeologic characteristics at a regional scale. Previous state-wide investigations that attempted to correlate atrazine sampling results with patterns of soils or geologic materials were unsuccessful and limited in approach. To better analyze the physical factors that might effect the distribution of atrazine detections, we decided to look at a smaller area, specifically Dane County.

Methods: In order to examine the physical controls on the distribution of atrazine detections, we had to determine the land area that contributes water to the sampled wells and then account for varying land-use practices within those areas. The project was conducted in four phases -- 1) development of GIS coverages of available soils, geologic, and hydrogeologic information; 2) determination of the zones of contribution (ZOCs) of wells sampled for atrazine; 3) estimation of atrazine application rates within the ZOCs and 4) statistical analysis to examine the relationships among the hydrogeologic factors, land-use patterns, and the detections of atrazine.

Results: Compilation of hydrogeologic data suggested the following results: 1)

Compiling domestic well-construction report information in a relational database and integrating the database with a GIS allow this information to be applied to a variety of problems. 2) PRETRAC, a pcARCPLLOT application developed for this project, greatly facilitated the development of input files for particle-tracking models as well as providing a powerful and flexible data exploration tool.

Statistical analysis of the relationships between atrazine detections and the variables describing physical and land-use factors were difficult to identify and quantify. Non-parametric techniques such as cross-tabulation of categorical data and logistic regression for multivariate analysis proved to be the most useful. Variables of primary importance in predicting whether a given domestic well would have an atrazine detection include 1) atrazine use, 2) presence of shale, 3) presence of Sinnipee Group dolomite as the uppermost bedrock unit over the majority of the ZOC, and 4) location in discharge area. Soil thickness, unlithified materials, and depth-to-bedrock data all suggest that thin, fine-grained soils may be attenuating atrazine.

Recommendations/

Implications: Characterizing soils, geology and understanding the groundwater flow system are important for prediction of atrazine detections. Land-use, presence of shale, and presence of Sinnipee Group dolomite were determined for the land area that contributes water to the well (or zone of contribution, ZOC, of the well). Flow system position, especially location in a discharge area also appears to be important. Fine-grained units, either shale, clay layers in the Sinnipee, or thin fine-grained soils all appear to attenuate atrazine.

The statistical model developed to predict atrazine detection using the detailed soils, geologic, and hydrogeologic data for all 325 wells had overall predictive accuracy of 66%. Stratification of the data and development of separate models for the glaciated and unglaciated portions of the county led to improved overall predictive accuracy. This suggests that prediction of the distribution of atrazine detections is complicated by the variability of natural settings and that predictive models developed for smaller areas, with more consistent soils, geologic, and hydrogeologic characteristics are more likely to be accurate.

Availability of Report:

The final report is available from the Wisconsin Geological and Natural History Survey, 3817 Mineral Point Road, Madison, WI, as Open File Report WOFR-94-02. A second open file report, WOFR 94-01, "Dane County Atrazine Project Database and Documentation: Programs, Data Sources and Methods" is available on diskette.

Related

Publications: Bohn, M.F., and Muldoon, M.A., 1993. Integrating Hydrogeologic Data and Models with a Geographic Information System: Applications from a Regional Pesticide Study. (Abstract) 1993 Environmental Systems Research Inc. User Conference.

Muldoon, M.A. and Bohn, M.F., 1993. Integrating Hydrogeologic Data from Well-construction Reports with a Geographic Information System: Applications from a Regional Pesticide Study. Proceedings of International Association of Hydrogeologists Workshop on the Development of Water Related Information Systems, Washington D.C., May 19-20, 1993, 9 p.

Key Words: Groundwater, atrazine, land-use, geographic information system (GIS)

Funding: The Wisconsin Department of Natural Resources provided funding for this project through the Groundwater Management Practice Monitoring Program which receives appropriations from the Groundwater Account. Additional funding was received from the Wisconsin Department of Agriculture, Trade, and Consumer Protection. The grant was matched from resources of the Dane County Land Conservation Department and the Wisconsin Geological and Natural History Survey who conducted this project cooperatively.