

Explanation

- p** **Sediment of low, typically wet areas.** Peat, muck, or slope sediment covering windblown, glacial, stream, or lake sediment as indicated by adjacent map units; low-lying, flat to low-relief surfaces.
- sp** **Postglacial stream sediment.** Silty and sandy sediment deposited by postglacial streams; flat to low-relief floodplains. Mapped where extensive; present but unmapped adjacent to most streams.
- ws** **Windblown sand.** Fine- to medium-grained sand; more than 1.5 m thick.
- su** **Meltwater-stream sediment, uncollapsed.** Gravelly sand deposited by meltwater streams; flat to low-relief surfaces; original depositional surfaces preserved in most places; covered in places by thin deposits of windblown sand (unit **suw**). May include areas of sandy sediment deposited in glacial lakes.
- sc** **Meltwater-stream sediment, collapsed.** Same as unit **su**, except that meltwater-stream sediment was deposited on ice; moderate to high-relief surfaces; original depositional surface destroyed by collapse; typically includes till and till-like debris-flow sediment where deposited adjacent to former ice margins; covered in places by thin deposits of windblown sand (unit **scw**).
- gn** **Glacial sediment of the Nashville Member of the Copper Falls Formation.** Reddish-brown to brown, sandy, typically non-dolomitic, slightly dolomitic in places, till and debris-flow sediment deposited by the Langlade Lobe in the far northwest part of the map area; found in areas of streamlined glacial topography.
- gsh** **Glacial sediment of the Silver Cliff Member of the Kewaunee Formation.** Reddish-brown, sandy, dolomitic till and debris-flow sediment; found in areas of moderate to high-relief hummocky topography (unit **gsh**); draped over Precambrian rock (unit **gsd**).
- gm** **Glacial sediment of the Middle Inlet Member of the Kewaunee Formation.** Reddish-brown, sandy, dolomitic till and debris-flow sediment in areas of rolling to low-relief hummocky topography (unit **gmh**); may include small areas of silty, dolomitic till of the Kirby Lake Member of the Kewaunee Formation.

UW
Extension

Wisconsin Geological and Natural History Survey
3817 Mineral Point Road, Madison, Wisconsin 53705-5100
608/263.7389 FAX 608/262.8086
www.uwex.edu/wgnhs/
James M. Robertson, Director and State Geologist

Digital cartography and editing by D.L. Patterson.

